

Available online at www.sciencedirect.com

SciVerse ScienceDirect

journal homepage: www.e-fjs.com

MINI-REVIEW

Reparative surgery for degenerative mitral valve regurgitation

Jen-Ping Chang*

*Division of Thoracic and Cardiovascular Surgery, Kaohsiung Chang Gung Memorial Hospital,
Chang Gung University College of Medicine, Kaohsiung, Taiwan*

Received 13 September 2011; received in revised form 26 September 2011; accepted 5 October 2011
Available online 6 December 2011

KEYWORDS

annuloplasty;
degenerative;
mitral regurgitation;
mitral valve repair

Summary Reparative surgery for degenerative mitral valve regurgitation or insufficiency has been the subject of countless articles in the literature over the past five decades. During this period, a wealth of experience has contributed to greater understanding of mitral valve anatomy and pathophysiology. Excellent surgical techniques for the treatment of mitral regurgitation have been developed and are undergoing further refinement. With the efforts of the pioneers, standard surgical techniques, strategies, and guidelines have been well established. These achievements significantly prolong the life-expectancy of patients, provide excellent long-term durability of the repaired valve, and reduce the risk of endocarditis. Mitral valve repair for degenerative mitral regurgitation also contributes to reduced risk of thromboembolic events and the need for life-long anticoagulation after mitral valve replacement. Copyright © 2011, Taiwan Surgical Association. Published by Elsevier Taiwan LLC. All rights reserved.

1. Introduction

Mitral valve repair is the best option for treatment of degenerative mitral regurgitation. This issue has been the subject of countless articles in the literature over the past

five decades. Prior reports have documented several advantages of mitral valve repair compared with replacement, including: lower operative mortality^{1,2}; better preservation of left ventricular function³; lower incidence of stroke and endocarditis^{4,5}; a lower rate of reoperation and complications of anticoagulation therapy; and superior long-term survival.⁶ In the United States, the Society of Thoracic Surgeons Adult Cardiac Surgery Database also documented the significantly lowered operative risk of mitral valve repair and the progressive adoption of this procedure from 51% to 69%.⁷ With the efforts of the pioneers, standard surgical techniques, strategies, and

* Corresponding author. Division of Thoracic and Cardiovascular Surgery, Kaohsiung Chang Gung Memorial Hospital, Chang Gung University College of Medicine, 123 Tapei Road, Niasung District, Kaohsiung City, Taiwan 833, ROC.
E-mail address: c9112772@adm.cgmh.org.tw.

guidelines have now been well established. The purpose of this mini review is to widen surgeons' acknowledgment of these important historic achievements, to summarize the modern state-of-the-art procedures, and to provide future perspectives on mitral valve repair for degenerative etiology.

2. Historical aspects

2.1. Restrictive annuloplasty

Direct attempts at mitral valve repair utilizing the techniques of suture annuloplasty under cardiopulmonary bypass support were initially described by Lillehei et al in 1957.⁸ With the exception of McGoon's technique of posterior leaflet plication,⁹ the mode of mitral valve repair was later modified by other surgeons with various types of commissural suture technique.^{10–12} However, by these palliative techniques, recurrence of mitral regurgitation or stenosis was not rare on account of leaflet or subvalvular apparatus dysfunction, persistent annular dilatation, and fibrosis of the plicated commissures.

2.2. Comprehensive remodeling concept

During the period from the 1970s to the 1980s, the concept of comprehensive reconstructive valvular surgery was introduced by Carpentier of Paris, the most widely recognized advocate of the annular remodeling concept, the so-called "French correction".^{13–15} Thus, the gate was opened to the modern comprehensive and sophisticated mitral reparative concept.

3. Modern state-of-the-art procedures

3.1. The role of ring annuloplasty

Mitral valve repair surgery has progressed and evolved dramatically since the inception of remodeling ring annuloplasty.^{13–16} The value of this particular method during concomitant reconstructive procedures has been firmly established. Mitral valve repair without an annuloplasty ring is an important factor influencing post-repair durability.¹⁷ Currently, ring annuloplasty is an essential component of mitral valve repair, as annular remodeling restores the anatomic shape of the annulus and ensures long-term maintenance of valve repair results.¹⁸

The rigid and semi-rigid rings proposed by Carpentier and his colleagues have been the most popular devices for remodeling ring annuloplasty.^{13–16} Meanwhile, Durán et al advocated a flexible ring concept for allowing a continuous change of the mitral annular conformation during the whole cardiac cycle.¹⁹ Furthermore, Chang et al extended this concept and proposed a pliable polytetrafluoroethylene (PTFE) partial ring for reducing the posterior annulus, with favorable results.²⁰ However, the debate regarding the ring rigidity ceased when Chang et al from Seoul found no significant clinical difference in the rate of recurrence of significant mitral regurgitation between rigid and flexible rings in a long-term follow-up study.²¹

Recently, saddle-shaped annuloplasty rings have been used increasingly more often to conform the mitral annulus to a nonplanar shape and possibly to reduce leaflet stress. The preliminary results seem promising, although larger and longer cohort studies are necessary.²²

3.2. Leaflet and chordal management

In a study of degenerative mitral valve repair, Gillinov et al reported the factors influencing repair durability: repair without an annuloplasty ring; repair without sliding annuloplasty; and the use of chordal shortening instead of chordal transposition or artificial chordal replacement.¹⁷ Since the introduction of PTFE as a chordal substitute, this has become the popular technique used by many surgeons instead of conventional procedures for correcting prolapse of the mitral leaflet, such as cusp resection, chordal shortening, and chordal transposition. This "respect but no resect" principle has changed the reparative technique for degenerative mitral regurgitation drastically and in most instances of degenerative mitral regurgitation repair, the conventional complex leaflet and subvalvular apparatus manipulation techniques have been aborted.^{23,24} The optimal length of artificial chordae has been obtained by several methods, but direct measurement of the nonprolapsed corresponding chordal height or measurement of distance of the papillary muscle tip to the corresponding annulus is recommended. In addition, the PTFE sutures are slippery and difficult to tie down at the intended length. Numerous methods have been described to ensure the correct length of the artificial chordae, however, most of these techniques are complex, requiring custom-made special devices, and time consuming.²⁵ Therefore, further refinement of this valuable technique is needed.

3.3. Edge-to-edge repair

The edge-to-edge (E2E) mitral valve repair was advocated in the late 1990s by Alfieri et al.²⁶ This technique was initially described as a rescue method for difficult or failed mitral valve repair with the standard technique. However, later data reported by the same research group confirmed overall suboptimal results of the E2E technique when annuloplasty is not added to the repair.²⁷ Currently, this technique has been preserved as an alternative option for prevention of the systolic anterior motion of the mitral anterior leaflet in repairing the severe form of myxomatous degenerative mitral regurgitation (Barlow's valve).

4. Future perspectives

4.1. Robotic minimally invasive valvular surgery

Robotic surgery was made feasible following the integration and improvements of telemanipulation, three-dimensional vision, the computer-enhanced EndoWrist, and a tremor-filtering system. Chitwood et al reported their experience in 300 cases of mitral repair with excellent results comparable with those of the conventional open technique.²⁸

However, few surgeons have adopted this state-of-the-art technology in their daily practice because the method is expensive, highly sophisticated, has a steep learning curve, and demands a meticulous combination of modern technologies.

4.2. Percutaneous catheter-based approach

Two-year follow-up data from the EVEREST II trial show that although a catheter-based mitral valve repair procedure using the MitraClip system was less effective in reducing mitral regurgitation than conventional surgery, similar improvements in clinical outcomes were observed with fewer short-term adverse events.²⁹ The major drawbacks of this catheter-based EZE technique are doubtful long-term durability and potential creation of a stenotic mitral valve.

Data from the EVOLUTION phase 1 study to assess the safety and efficacy of the percutaneous coronary sinus device (MONARC device) show limited feasibility and efficacy. It is noteworthy that coronary artery compression may occur in patients whose greater cardiac vein passes over a coronary artery, necessitating some future strategies to avoid this serious complication.³⁰

The development of robotic surgery and the catheter-based approaches for repairing mitral regurgitation are important and inevitable trends. At present, however, considering the current excellent long-term survival and results achieved from the conventional open reparative techniques for degenerative mitral regurgitation, a very cautious attitude towards application and evaluation of such new technologies is mandatory.

5. Summary

Mitral valve repair for degenerative mitral regurgitation is a well-established therapeutic option and has been practiced worldwide. Unfortunately, the reported number of cases using this procedure in Taiwan remains smaller than that reported in North America and European countries.⁷ Although mitral valve repair is technically demanding, with the accumulation of sufficient experience, success should be achieved with consistency. We, the cardiac surgeons practicing in the current era, should consider every patient with degenerative mitral regurgitation, make all efforts to find reparable cases, and attempt repair in all suitable cases.

References

- Enriquez-Sarano M, Schaff HV, Orszulak TA, Tajik AJ, Bailey KR, Frye RL. Valve repair improves the outcome of surgery for mitral regurgitation. A multivariate analysis. *Circulation*. 1995; 91:1022–1028.
- Rankin JS, Hammill BG, Ferguson TB, et al. Determinants of operative mortality in valvular heart surgery. *J Thorac Cardiovasc Surg*. 2006;131:547–557.
- David TE, Burns RJ, Bacchus CM, Druck MN. Mitral valve replacement for mitral regurgitation with and without preservation of chordae tendineae. *J Thorac Cardiovasc Surg*. 1984;88:718–725.
- Braunberger E, Deloche A, Berrebi A, et al. Very long-term results (more than 20 years) of valve repair with Carpentier's techniques in nonrheumatic mitral valve insufficiency. *Circulation*. 2011;124(12 Suppl 1):18–11.
- Gillinov AM, Faber CN, Sabik JF, et al. Endocarditis after mitral valve repair. *Ann Thorac Surg*. 2002;73:1813–1816.
- Jokinen JJ, Hippeläinen MJ, Pitkänen OA, Hartikainen JE. Mitral valve replacement versus repair: propensity-adjusted survival and quality-of-life analysis. *Ann Thorac Surg*. 2007; 84:451–458.
- Gammie JS, Sheng S, Griffith BP, et al. Trends in mitral valve surgery in the United States: results from the society of thoracic surgeons adult cardiac surgery database. *Ann Thorac Surg*. 2009;87:1431–1439.
- Lillehei CW, Gott VL, DeWall RA, Varco RL. Surgical correction of pure mitral insufficiency by annuloplasty under direct vision. *J Lancet*. 1957;2:446–449.
- McGoon DC. Repair of mitral insufficiency due to ruptured chordae tendineae. *J Thorac Cardiovasc Surg*. 1960;39:357–362.
- Wooler GH, Nixon PG, Crimshaw VA, Watson DA. Experience with the repair of the mitral valve in mitral incompetence. *Thorax*. 1962;17:49–57.
- Kay JH, Egerton WS, Zubieta P. The surgical treatment of mitral insufficiency and combined mitral stenosis and insufficiency with use of the heart-lung machine. *Surgery*. 1961;50: 67–74.
- Reed GE, Tice DA, Clauss RH. Asymmetric exaggerated mitral annuloplasty: repair of mitral insufficiency with hemodynamic predictability. *J Thorac Cardiovasc Surg*. 1965;49: 752–761.
- Carpentier A, Relland J, Deloche A, et al. Conservative management of the prolapsed mitral valve. *Ann Thorac Surg*. 1978;26:294–302.
- Carpentier A, Chauvaud S, Fabiani JN, et al. Reconstructive surgery of mitral valve incompetence: ten-year appraisal. *J Thorac Cardiovasc Surg*. 1980;79:338–348.
- Carpentier A. Cardiac valve surgery: The "French correction". *J Thorac Cardiovasc Surg*. 1983;86:323–337.
- Carpentier AF, Lessana A, Relland JY, et al. The "Physio-Ring": an advanced concept in mitral valve annuloplasty. *Ann Thorac Surg*. 1995;60:1177–1185.
- Gillinov AM, Cosgrove DM, Blackstone EH, et al. Durability of mitral valve repair for degenerative disease. *J Thorac Cardiovasc Surg*. 1998;116:734–743.
- Schwartz CF, Gulkarov I, Bohmann K, Colvin SB, Galloway AC. The role of annuloplasty in mitral valve repair. *J Cardiovasc Surg*. 2004;45:419–425.
- Durán CM, Pomar JL, Cucchiara G. A flexible ring for atrio-ventricular heart valve reconstruction. *J Cardiovasc Surg (Torino)*. 1978;19:417–420.
- Chang CH, Lin PJ, Chang JP, Chu JJ, Hsieh MJ, Chiang CW. Long-term results of polytetrafluoroethylene mitral annuloplasty. *Ann Thorac Surg*. 1994;57:644–647.
- Chang BC, Youn YN, Ha JW, Lim SH, Hong YS, Chung N. Long-term clinical results of mitral valvuloplasty using flexible and rigid rings: A prospective and randomized study. *J Thorac Cardiovasc Surg*. 2007;133:995–1003.
- Mahmood F, Gorman JH 3rd, Subramaniam B, et al. Changes in mitral valve annular geometry after repair: saddle-shaped versus flat annuloplasty rings. *Ann Thorac Surg*. 2010;90: 1212–1220.
- Kasegawa H, Shimokawa T, Shibasaki I, Hayashi H, Koyanagi T, Ida T. Mitral valve repair for anterior leaflet prolapse with expanded polytetrafluoroethylene sutures. *Ann Thorac Surg*. 2006;81:1625–1631.

24. Kuntze T, Borger MA, Falk V, et al. Early and mid-term results of mitral valve repair using premeasured Gore-Tex loops ("loop technique"). *Eur J Cardiothorac Surg*. 2008;33: 566–572.
25. Chang JP, Kao CL. Slit stent technique for ensuring the correct length of artificial chordae in mitral repair. *J Card Surg*. 2011;26: 259–260.
26. Alfieri O, Maisano F, De Bonis M, et al. The double-orifice technique in mitral valve repair: A simple solution for complex problems. *J Thorac Cardiovasc Surg*. 2001;122:674–681.
27. Maisano F, Caldarola A, Blasio A, De Bonis M, La Canna G, Alfieri O. Midterm results of edge-to-edge mitral valve repair without annuloplasty. *J Thorac Cardiovasc Surg*. 2003;126:1987–1997.
28. Chitwood WR Jr, Rodriguez E, Chu MW, et al. Robotic mitral valve repairs in 300 patients: A single-center experience. *J Thorac Cardiovasc Surg*. 2008;136:436–441.
29. Cleland JG, Coletta AP, Freemantle N, Clark AL. Clinical trials update from the American College of Cardiology Meeting 2011: STICH, North Star, TARGET, and EVEREST II. *Eur J Heart Fail*. 2011;13:805–808.
30. Harnek J, Webb JG, Kuck KH, et al. Transcatheter implantation of the MONARC coronary sinus device for mitral regurgitation: 1-year results from the EVOLUTION phase 1 study (Clinical evaluation of the Edwards lifesciences percutaneous mitral annuloplasty system for the treatment of mitral regurgitation). *JACC Cardiovasc Interv*. 2011;4:115–122.